

ORIGINAL PAPER

## High resilience of coral communities at Hin Ang Wang, Surat Thani Province and a proposed management strategy

Sittiporn Pengsakun<sup>a</sup>, Supawadee Hamanee<sup>b</sup>, Oranit Saengthongsuk<sup>b</sup>, Anupong Avirutha<sup>c</sup>, Panida Rakklin<sup>c</sup>, Wanlaya Klinthong<sup>a</sup>, Thamasak Yeemin<sup>a</sup>, Makamas Sutthacheep<sup>a,\*</sup>

<sup>a</sup>Marine Biodiversity Research Group, Department of Biology, Faculty of Science, Ramkhamhaeng University, Huamark, Bangkok, Thailand

<sup>b</sup>School of Business Administration, Sripatum University, Jatujak, Bangkok

<sup>c</sup>Collage of Tourism and Hospitality, Sripatum University, Jatujak, Bangkok

\*Corresponding author: msutthacheep@yahoo.com

Received: 20 August 2022/ Revised: 26 August 2022/ Accepted: 28 August 2022

**Abstract** The 1998 and 2010 mass coral bleaching events caused coral degradation to be severe and extensive in the Gulf of Thailand, particularly the Western Gulf of Thailand. This study assessed the resilience potential of coral communities at Hin Ang Wang, the Western Gulf of Thailand, by conducting field surveys on benthic components, community structure of scleractinian corals, diversity and density of juvenile corals, composition and abundance of macrobenthic invertebrates and reef fish. The live coral cover was 44.2%, while dead coral cover was 3.7%, indicating that the coral community is very good condition. The dominant coral species were *Porites lutea*, *Pavona decussata* and *Montipora aequituberculata*. The total density of juvenile corals was 1.6 juveniles/m<sup>2</sup>. The total density of 27 reef fish species was 937. 8±433.6 individuals/100 m<sup>2</sup>. The most dominant reef fishes were *Neopomacentrus anabatooides*, *Chromis cineracens* and *Parioglossus philippinus*. There were several target species of macrobenthic invertebrates and reef fish for aquaculture and fishing. Some are attractive to tourists and can be target species for marine ecotourism. Our results show that the coral communities at Hin Ang Wang had high resilience to coral bleaching events and human impacts. It should be established as a marine protected area under the Department of Marine and Coastal Resources. The management plan should be also developed and implemented. The resilience-based management is proposed to support natural processes and maintain coral reef resilience

**Keywords:** coral community, resilience, management, recruitment, Gulf of Thailand

### 1. Introduction

Coral reefs worldwide are degrading because of human and natural disturbances, especially land-based pollution, overfishing, ocean warming and ocean acidification (Yeemin et al. 2009; Yeemin et al. 2013a; Wear and Thurber 2015; Cheal et al. 2017; Hughes et al. 2017; Shantz et al. 2020). The future of coral reef ecosystems relies on the collaborative reduction of global greenhouse gas emissions and urgent actions that enhance coral reef resilience to climate change, particularly coral bleaching events (Kleypas et al. 2021; Shaver et al. 2021). The severe coral bleaching events have caused the majority of coral reefs in loss of living corals around the world (Eakin et al. 2019). Some coral reefs are vulnerable to collapse at the regional level and some ecoregions are critically endangered (Obura et al. 2022).

The 1998 and 2010 mass coral bleaching events caused coral degradation to be severe and extensive in the Gulf of Thailand, particularly the Western Gulf of Thailand. Scleractinian corals in the Gulf of Thailand are different in their susceptibility to coral bleaching, which is an essential aspect of coral reef research because it can significantly lead to coral community changes

(Yeemin et al. 2012; Sutthacheep et al. 2012). Knowledge concerning the synergistic effects of coral bleaching and anthropogenic impacts on the ecological functions and processes of coral reefs, especially coral recruitment, is very important for establishing a management strategy for enhancing the resilience of coral reefs (Manikandan et al. 2017). High coral recruitment and survivorship of juvenile corals are essential factors in maintaining coral populations after mass mortality from bleaching events (Hughes et al. 2002; Doropoulos et al. 2015; Yucharoen et al. 2015). Therefore, coral recruitment is usually applied as a bioindicator of coral reef recovery and resilience after severe coral bleaching events. Several environmental factors influence coral recruitment, particularly sedimentation, water pollution and overfishing (Yeemin et al. 2013b).


The capacity of coral reefs to resist or recover from degradation and to maintain their ecosystem services are defined as coral reef resilience (Mumby et al. 2007). Resilience-based management of coral reefs assesses spatial variation in resilience potential and implementing proper management plans. Several resilience indicators have been developed and used for assessing the ecological resilience of coral reefs, including the assessment of the resilience potential of inshore and offshore coral communities in the Western Gulf of Thailand (Sutthacheep et al. 2019). This study quantitatively assessed the coral reef resilience potential at Hin Ang Wang, Surat Province, the Western Gulf of Thailand, based on scientific surveys in early 2022.

## 2. Material and Methods

### 2.1 Study site

The study site is located at Hin Ang Wang (9°23'27.80"N, 100° 1'31.01"E), Samui Island,

Surat Thani Province, the Western Gulf of Thailand. The coral reef area is approximately 78,000 square meters.



**Figure 1.** Map of the study site at Hin Ang Wang, Surat Thani Province, the Western Gulf of Thailand

### 2.2 Data collection and analysis

The study was conducted on coral communities at Hin Ang Wang in early 2022. The coral communities were found at approximately 0-10 m in depth. Live coral cover was recorded in three permanent belt-transects of 30x1 m<sup>2</sup> and scleractinian corals (>5 cm diameter) were identified to species level. The number of visible juvenile corals (1-5 cm diameter) was also measured in these three permanent belt-transects. All juvenile corals were identified to genus. (Veron 2000).

The numbers of each macrobenthic invertebrate species found on the coral communities were recorded and counted in three permanent belt-transects of 50x1 m<sup>2</sup> and they were identified to a species level, if possible.

Coral reef fishes were observed using a standard visual census technique (English et al. 1997). The belt transect, 50x5 m was conducted at each study site. SCUBA divers estimated fish abundance and identified species of reef fish along each transect. Species not recognized in the field surveys were photographed and video-recorded for identification in the laboratory. Fishes were identified to species level, if possible.


### 3. Results

The benthic components investigation revealed that live coral and rock were the major components of the substrates at Hin Ang Wang (Figure 2). The other components were sand, rubble, dead coral and other sessile invertebrates, including soft corals


and sea anemones (Figure 3). The live coral cover was 44.2%, while dead coral cover was 3.7%, indicating that the coral community is very good condition. The dominant coral species were *Porites lutea*, *Pavona decussata* and *Montipora aequituberculata*. Many large colonies of *Diploastrea heliopora* and *Porites lutea* were also found at the deeper zone. The abundant coral species were *Acropora hyacinthus*, *Favites abdita*, *Galaxea astreata*, *Galaxea fascicularis*, *Goniopora columna*, *Lobophyllia hemprichii*, *Lobophyllia radians*, *Merulina ampliata*, *Pachyseris speciosa*, *Pavona cactus*, *Pocillopora acuta*, *Podabacia crustacea*, *Pectinia lactuca*, *Herpolitha limax* and *Turbinaria peltata*. (Figures 4 and 5).



**Figure 2.** The coral reef community at Hin Ang Wang, Surat Thani Province, the Western Gulf of Thailand



**Figure 3.** Benthic components at the study site (mean ± SD)



**Figure 4.** Species composition of live coral cover at the study site (mean ± SD)



*Porites lutea*

*Pavona decussata*

*Montipora aequituberculata*

**Figure 5.** The dominant coral species at the study site



The total density of juvenile corals at Hin Ang Wang was 1.6 juveniles/m<sup>2</sup>. The most abundant juvenile coral was *Fungia*. The common juvenile corals were *Acropora*,

*Ctenactis*, *Dipsastraea*, *Echinopora*, *Favites*, *Hydnophora*, *Litthophyllon*, *Montipora*, *Pavona*, *Platygyra*, *Pocillopora* and *Porites* (Figure 6).




Figure 6. Density of juvenile corals on available substrates at the study site (mean ± SD)

The most dominant macrobenthic invertebrate at Hin Ang Wang was the Christmas tree worms, *Spirobranchus giganteus* (5.4 individuals/m<sup>2</sup>). The other macrobenthic invertebrates were *Neopetrosia* sp., *Xestospongia* sp., *Rochia nilotica*, *Drupella rugosa*, *Phyllidia* sp., *Arca* sp.,

*Begonia semiorbiculata*, *Tridacna crocea*, *T. squamosa*, *Hyotisa hyotis*, *Chama* sp., *Pedum spondyloidum*, *Spondylus* sp. and *Diadema setosum*. Some macrobenthic invertebrates are attractive to tourists, such as *S. giganteus* and *T. squamosa* (Figures 7 and 8).




Figure 7. Abundance of macrobenthic invertebrate at the study site (mean ± SD)



*Spirobranchus giganteus*




*Tridacna squamosa*

**Figure 8.** Some attractive macrobenthic invertebrates to tourists.

The densities of reef fish at Hin Ang Wang are shown in Figure 9. The total density of 27 reef fish species was  $937.8 \pm 433.6$  individuals/100 m<sup>2</sup>. The most dominant reef fishes were *Neopomacentrus anabatooides*, *Chromis cineracens* and *Parioglossus philippinus*.

The fishing target was *Lutjanus carponotatus*, *Lutjanus russellii*, *Caesio*

*cuning*, *Plectorhinchus gibbosus*, *Scarus rivulatus* and *Cephalopholis boenak*. Some reef fish are attractive to tourists, including *Sargocentron rubrum*, *Ostorhinchus endekataenia*, *Parioglossus philippinus*, *Chaetodon octofasciatus* and *Chelmon rostratus* (Figure 10).



**Figure 9.** Species composition of reef fish at the study site (mean  $\pm$  SD)



*Chelmon rostratus*



*Sargocentron rubrum*

**Figure 10.** Some attractive reef fish to tourists

#### 4. Discussion

The community structure of scleractinian corals at Hin Ang Wang observed from this study showed high resilience after the severe coral bleaching events in 1998 and 2010. The high percentage of live coral cover and low percentages of dead coral and rubble were obviously recorded. The high susceptibility of *Pocillopora acuta*, *Montipora* spp., *Acropora* spp. and *Turbinaria* spp., in the Gulf of Thailand (Sutthacheep et al. 2010; Yeemin et al. 2010; Sutthacheep et al. 2013) were found at Hin Ang Wang. Some tolerant corals to the coral bleaching events, such as *Galaxea*, *Pachyseris*, *Turbinaria*, *Favites* and *Diploastrea* were abundantly recorded at Hin Ang Wang.

Our results show juvenile corals from many species, particularly *Acropora*, *Ctenactis*, *Dipsastraea*, *Echinopora*, *Favites*, *Fungia*, *Hydnophora*, *Litthophyllon*, *Montipora*, *Pavona*, *Platygyra*, *Pocillopora*, *Porites* were observed at Hin Ang Wang. These data imply the connectivity of some coral populations in the Western Gulf of Thailand. Further studies using molecular genetic techniques would confirm the coral genetic connectivity and self-seeding in this region

(van Oppen et al. 2008; Klinthong et al. 2019). The density of juvenile corals reported in this study was 1.6 juveniles/m<sup>2</sup>, which is comparable to previous reports in the Biscayne National Park (Miller et al. 2000), Mesoamerican barrier reef (Ruiz-Zarate and Arias Gonzalez 2004), and Indian Ocean reef (Manikandan et al. 2017).

Major macrobenthic invertebrates found at Hin Ang Wang are categorized into 5 classes, namely Demospongiae, Polychaeta, Gastropoda, Bivalvia and Echinoidea. Several species of macrobenthic invertebrates are from the Bivalvia, namely *Arca* sp., *Begonia semiorbiculata*, *Tridacna crocea*, *T. squamosa*, *Hytisa hyotis*, *Chama* sp., *Pedum spondyloidum* and *Spondylus* sp. These bivalves have high potential to be commercial species in mariculture, particularly the giant clams *Tridacna* spp. The giant clam mariculture industry has been providing food and marine aquarium trade markets for over four decades (Militz et al. 2017). Research on artificial interspecific hybridization of two giant clams, *T. squamosa* and *T. crocea*, in the South China Sea revealed that the cross between *T. squamosa* and *T. crocea* was successfully carried out. The reciprocal

hybrids showed larger growth heterosis, survival advantage and a beautiful mantle coloration. These hybrids exhibited high potential for application in the market and giant clam mariculture industry (Zhou et al. 2020). Our results also imply several target species of macrobenthic invertebrates and reef fish at Hin Ang Wang for aquaculture and fishing, especially *T. crocea*, *T. squamosa*, *Diadema setosum*, *Lutjanus carponotatus*, *L. russellii*, *Caesio cuning*, *Plectorhinchus gibbosus*, *Scarus rivulatus* and *Cephalopholis boenak*. Moreover, several species of macrobenthic invertebrates and reef fish at Hin Ang Wang, such as *Neopetrosia* sp., *Xestospongia* sp., *Spirobranchus giganteus*, *Phyllidia* sp., *Tridacna* spp., *Pedum spondyloidum*, *Sargocentron rubrum*, *Ostorhinchus endekataenia*, *Parioglossus philippinus*, *Chaetodon octofasciatus* and *Chelmon rostratus*, are attractive to tourists and can be target species for marine ecotourism (Wongnutpranont et al. 2020; Rangseethampanya et al. 2021; Yeemin et al. 2021).

Coral reefs can be managed by using several management tools. Recently, resilience-based management (RBM) has been applied coral reef management with concentrating on the use of knowledge of existing and possible drivers that potentially influence the functions and structure of coral reef ecosystem, such as climate change, coral disease outbreaks, land-use change, coastal development and reef fisheries (McLeod et al. 2019; Sutthacheep et al. 2022). Our results show that the coral communities at Hin Ang Wang had high resilience to coral bleaching events and human impacts. Hin Ang Wang should be established as a marine protected area under the Department of Marine and Coastal Resources to protect the healthy coral community. Our results also imply that Hin Ang Wang should implement its

management plans properly to maintain its resilience. RBM may be used to support natural processes. Marine ecotourism should be promoted as it can protect the coral community and keep the tourist numbers below the carrying capacity of this tourist destination (Sutthacheep et al. 2019).

### Acknowledgements

We are most grateful to the staff of Marine Biodiversity Research Group, Faculty of Science, Ramkhamhaeng University, for their support and assistance in the field. This work was supported by Ramkhamhaeng University (RU), Thailand Science Research and Innovation (TSRI) and National Science, Research and Innovation Fund (NSRF).

### References

- Cheal AJ, MacNeil MA, Emslie MJ, Sweatman H (2017) The threat to coral reefs from more intense cyclones under climate change. *Glob Change Biol* 23(4):1511–1524. <https://doi.org/10.1111/gcb.13593>
- Doropoulos C, Ward S, Ro G, González-Rivero M, Mumby PJ (2015) Linking demographic processes of juvenile corals to benthic recovery trajectories in two common reef habitats. *PLoS ONE* 10: e0128535
- Eakin CM, Sweatman H, Brainard RE (2019) The 2014–2017 global-scale coral bleaching event: Insights and impacts. *Coral Reefs* 38(4): 539–545. <https://doi.org/10.1007/s00338-019-01844-2>
- English S, Wilkinson C, Baker V (1997). Survey manual for tropical marine resources. Australian Institute of Marine Science. 390 pp.
- Hughes TP, Baird AH, Dinsdale EA, Harriott VJ, Moltschaniwskyj NA, Pratchett MS, Tanner JE, Willis BL (2002) Detecting regional variation



- using meta-analysis and large-scale sampling: Latitudinal patterns in recruitment. *Ecology* 83: 436–451
- Hughes TP, Kerry JT, Álvarez-Noriega M, et al (2017) Global warming and recurrent mass bleaching of corals. *Nature* 543(7645): 373–377. <https://doi.org/10.1038/nature21707>
- Kleypas J, Allemand D, Anthony K, Baker AC, Beck MW, Hale LZ, Hilmi N, Hoegh-Guldberg O, Hughes T, Kaufman L, Kayanne H, Magnank AK, Mcleod E, Mumby P, Palumbi S, Richmond RH, Rinkevich B, Steneck RS, Voolstra CR, Gattuso JP (2021). Designing a blueprint for coral reef survival. *Biol. Conserv* 257:109107. <https://doi.org/10.1016/j.biocon.2021.109107>
- Klinthong W, Pengsakun S, Sutthacheep M, Ruangthong C, Chamchoy C, Yeemin T (2019) Recovery of corals in Mu Ko Ang Thong, the Western Gulf of Thailand. *RIST* 2(1): 8-16
- Manikandan B, Ravindran J, Vidya PJ, Shrinivasu S, Manimurali R, Paramasivam K (2017) Resilience potential of an Indian Ocean reef: an assessment through coral recruitment pattern and survivability of juvenile corals to recurrent stress events. *Environ Sci Pollut Res* 24: 3614-13625
- Mcleod E, Anthony KRN, Mumby PJ, Maynard J, Beedene R, Graham NAJ et al (2019) The future of resilience-based management in coral reef ecosystems. *J Environ Manag* 233:291–301
- Miller MW, Weil E, Szmant AM (2000) Coral recruitment and juvenile mortality as structuring factors for the reef benthic communities in Biscayne National Park, USA. *Coral Reefs* 19:115–123
- Militz TA, Foale S, Kinch J, Southgate PC (2017) Consumer perspectives on theoretical certification schemes for the marine aquarium trade. *Fish Res* 193: 33-42
- Mumby PJ, Hastings A, Edwards JG (2007) Thresholds and the resilience of Caribbean coral reefs. *Nature* 450: 98–101
- Obura, D., Gudka, M., Samoilys, M. et al. (2022). Vulnerability to collapse of coral reef ecosystems in the Western Indian Ocean. *Nat Sustain* 5:104–113 <https://doi.org/10.1038/s41893-021-00817-0>
- Rangseethampanya P, Sutthacheep M, Ruangthong C, Yeemin T (2021) Distribution of *Chaetodon wiebeli*, a common ornamental fish, in Mu Ko Chumphon National Park. *RIST* 4(3): 52-59
- Ruiz-Zarate MA, Arias Gonzalez JE (2004) Spatial study of juvenile corals in the northern region of the Mesoamerican Barrier Reef System (MBRS). *Coral Reefs* 23:584–594
- Shantz AA, Ladd MC, Burkepile DE (2020). Overfishing and the ecological impacts of extirpating large parrotfish from Caribbean coral reefs. *Ecol Monogr* 90(2): e01403. <https://doi.org/10.1002/ecm.1403>
- Shaver EC, McLeod E, Hein MY, Palumbi SR, Quigley K, Vardi T, Mumby PJ, Smith D, Montoya-Maya P, Muller EM (2022) A roadmap to integrating resilience into the practice of coral reef restoration. *Glob Chang Biol* 28: 4751–4764
- Sutthacheep M, Chamchoy C, Pengsakun S, Klinthong W, Yeemin T (2019) Assessing the resilience potential of inshore and offshore coral communities in the Western Gulf of

- Thailand. *J Mar Sci Eng* 7:408.  
<https://doi.org/10.3390/jmse7110408>
- Sutthacheep M, Pongsakun S, Yucharoen M, Klinthong W, Sangmanee K, Yeemin T (2013) Impacts of the mass coral bleaching events in 1998 and 2010 on the western Gulf of Thailand. *Deep-Sea Research II* 96: 25-31
- Sutthacheep M, Saenghaisuk C, Pongsakun S, Yeemin T (2010) Bleaching and mortality of scleractinian corals at Hin Rap, Trat Province. In *Proceedings of 36<sup>th</sup> Congress on Science and Technology of Thailand: 26 – 28 October 2010, Bangkok, Thailand*. 4 pp.
- Sutthacheep M, Yeemin T, Aliño PM (2022). Reef Ecology in the Western Pacific for Adaptation to Global Change. In: Zhang J, Yeemin T, Morrison RJ, Hong GH (eds) *Coral Reefs of the Western Pacific Ocean in a Changing Anthropocene. Coral Reefs of the World, vol 14*. Springer, Cham. [https://doi.org/10.1007/978-3-030-97189-2\\_4](https://doi.org/10.1007/978-3-030-97189-2_4)
- Sutthacheep M, Yucharoen M, Klinthong W, Pongsakun S, Sangmanee K, Yeemin T (2012) Coral mortality following the 2010 mass bleaching event at Kut Island, Thailand. *Phuket Mar Biol Cent Res Bull* 71: 83–92
- van Oppen MJH, Lutz A, De'ath G, Peplow L, Kininmonth S (2008) Genetic traces of recent long-distance dispersal in a predominantly self-recruiting coral. *PLoS One* 3:e3401
- Veron JEN (2000) *Corals of the World*. Vol. 1–3. Australian Institute of Marine Science and CRR, Queensland, Australia
- Wear SL, Thurber RV (2015). Sewage pollution: Mitigation is key for coral reef stewardship. *Ann N Y Acad Sci* 1355(1): 15–30. <https://doi.org/10.1111/nyas.12785>
- Wongnutpranont A, Pongsakun S, Ruengthong C, Hamanee S, Sutthacheep M, Yeemin T (2020) Assessing potential sites for marine ecotourism in Chumphon Province, Thailand. *Ramkhamhaeng International Journal of Science and Technology* 3(3): 21-29
- Yeemin T, Pongsakun S, Yucharoen, M., Klinthong, W., Sangmanee, K., Sutthacheep, M. (2013a). Long-term decline in *Acropora* species at Kut Island, Thailand, in relation to coral bleaching events. *Marine Biodiversity* 43:23–29
- Yeemin T, Pongsakun S, Yucharoen M, Klinthong W, Sangmanee K, Sutthacheep M (2013b) Long-term changes of coral communities under stress from sediment. *Deep-Sea Research II* 96: 32-40
- Yeemin T, Saenghaisuk C, Sutthacheep M, Pongsakun S, Klinthong W, Sangmanee K (2009) Conditions of coral communities in the Gulf of Thailand: a decade after the 1998 severe bleaching event. *Galaxea* 11: 207-217
- Yeemin T, Saenghaisuk C, Yucharoen M, Klinthong W, Sutthacheep, M (2012) Impact of the 2010 coral bleaching event on survival of juvenile coral colonies in the Similan Islands, on the Andaman Sea coast of Thailand. *Phuket Mar Biol Cent Res Bull* 71: 93–102
- Yeemin, T, Sutthacheep M, Klinthong W, Sangmanee N, Chamchoy C, Jungrak L (2021) Abundance of the magnificent sea anemone (*Heteractis magnifica*) and its marine ecotourism potential at Mu Ko Chumphon

National Park, Thailand. RIST 4(2):  
11-18

Yucharoen M, Yeemin T, Casareto BE,  
Suzuki Y, Samsuvan W, Sangmanee  
K, Klinthong W, Pongsakun S,  
Sutthacheep M (2015) Abundance,  
composition and growth rate of coral  
recruits on dead corals following the  
2010 bleaching event at Mu Ko Surin,  
the Andaman Sea. Ocean Sci J 50:  
307–315

Zhou Z, Li J, Ma H, Qin Y, Zhou Y, Wei J,  
et al. (2020) Artificial interspecific  
hybridization of two giant clams,  
*Tridacna squamosa* and *Tridacna*  
*crocea*, in the South China Sea.  
Aquaculture, 515: 734581